IMMUNE NETWORK

Editorial

(Check for updates

Back to the T Cell: Basic and Clinical Application

Yong Woo Jung 💿 ^{1,*}, Su-Hyung Park 💿 ^{2,*}, Chang-Duk Jun 💿 ^{3,*}

¹Department of Pharmacy, College of Pharmacy, Korea University, Sejong 30019, Korea ²Graduate School of Medical Science and Engineering, Korea Advanced Institute of Science and Technology (KAIST), Daejeon 34141, Korea

³Immune Synapse and Cell Therapy Research Center, School of Life Sciences, Gwangju Institute of Science and Technology (GIST), Gwangju 61005, Korea

Since the first vaccine was tried in human medicine in the 1700s, immunity has been a central focus for protecting our health and lives. However, the key players in the immune system were not discovered until the 1960s, when Good and his colleagues identified 2 types of lymphocytes in chickens (1). These groundbreaking findings opened up a new field in immunology, but it took another decade for Zinkernagel and Doherty (2) to determine how T cells recognize antigens. Since then, much progress has been made in understanding the fundamental functions of T cells in health and disease.

In addition to these foundational research findings, there have been significant advances in the clinical implications of T cells. The most prominent T cell-based immunotherapies include, but are not limited to, immune checkpoint therapy, chimeric antigen receptor-T cell therapy, and vaccines (3). Other immunotherapies indirectly or directly modify T cells for therapeutic purposes. Since the current coronavirus disease 2019 pandemic, interest in T cell responses to disease has never been higher. In particular, the importance of vaccine or infection-induced T cell responses in controlling viral infections has been increasingly emphasized, especially after reports of neutralizing antibodies against severe acute respiratory syndrome coronavirus 2 variants. These recent developments have encouraged us to re-evaluate T cells to gain a deeper understanding of their precise roles in these immunotherapies and to engineer T cells to achieve better therapeutic outcomes.

In this special issue of the Immune Network, a collection of review articles provides a comprehensive overview of T cells and their regulation. It begins by examining the heterogeneity of naïve T cells, which is crucial for T cells to respond to a wide range of antigens. This review article presents the current knowledge on how the diversity of naïve T cells can lead to the formation of various effector and memory subpopulations in response to antigenic stimulation, and how the fate of T cell responses is determined (4). Additionally, a manuscript explores recent findings on T cell microvilli, which can release TCR after interacting with antigen-presenting cells, and their role in regulating immune responses to antigens (5).

The development of effector and memory T cells is vastly different between CD4⁺ and CD8⁺ T cells. While CD4⁺ Th cells can be classified into various types of subpopulations based on the cytokines they produce, the diversity of effector CD8⁺ CTLs is limited. Since the first description of the Th1/Th2 paradigm, many different subsets of Th cells have been discovered, and the key factors influencing their formation and functions, including cytokines and transcription factors, have been intensively studied. The current

OPEN ACCESS

Published online: Feb 24, 2023

*Correspondence to

Yong Woo Jung

Department of Pharmacy, College of Pharmacy, Korea University, 2511 Sejong-ro, Sejong 30019, Korea. Email: yjung@korea.ac.kr

Su-Hyung Park

Graduate School of Medical Science and Engineering, Korea Advanced Institute of Science and Technology (KAIST), 291 Daehakro, Yuseong-gu, Daejeon 34141, Korea. Email: park3@kaist.ac.kr

Chang-Duk Jun

Immune Synapse and Cell Therapy Research Center, School of Life Sciences, Gwangju Institute of Science and Technology (GIST), 123 Cheomdangwagi-ro, Buk-gu, Gwangju 61005, Korea.

Email: cdjun@gist.ac.kr

$\ensuremath{\textbf{Copyright}}\xspace$ © 2023. The Korean Association of Immunologists

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (https:// creativecommons.org/licenses/by-nc/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Yong Woo Jung D https://orcid.org/0000-0002-5599-0553 Su-Hyung Park D https://orcid.org/0000-0001-6363-7736 Chang-Duk Jun D https://orcid.org/0000-0002-1372-9277

1/3

Generated by 🛟 xmlinkpress

IMMUNE NETWORK

Conflict of Interest

The authors declare no potential conflicts of interest.

Author Contributions

Conceptualization: Jung YW, Park SH, Jun CD; Supervision: Jun CD; Writing - original draft: Jung YW, Park SH, Jun CD. understanding of these factors is described in a review article (6). Other crucial factors for metabolic and epigenetic regulation and fate decision, as well as factors related to the plasticity and heterogeneity of Th cells, are also summarized (7). Among these subsets, the specific plasticity between Th17 and Treg has been a point of interest in the gut, primarily due to the factors produced by microorganisms in the organ (8). These microorganisms have also received attention because they produce metabolites that modulate the differentiation and functions of Th cells (9).

CD8⁺ T cells have also attracted much attention for their potential to kill abnormal cells in immunotherapies for tumors and viral infections. CTLs are a heterogeneous population, and it is crucial to understand the characteristics of each subpopulation for clinical implications. One review article focuses on the regulation of CTL function based on the expression of NK receptors, which determines the functions and regulations of CTL subsets (10). Another article highlights the importance of metabolic switch in the tumor microenvironment for regulating CTL function, as nutrients and critical metabolites are limited (11).

Small subsets of effector T cells differentiate into memory T (Tm) cells after clearing the antigen, which are long-lived and respond robustly to the same antigens. However, Tm cells are also heterogeneous, so they must be carefully selected for application in human diseases (12). Another recent discovery, the "memory-like" T cell population called virtual memory T (Tvm) cells, which acquire a memory phenotype in the absence of foreign Ag, is reviewed. The article outlines the distinct characteristics, immunological properties, and potential application of Tvm cells in immunotherapy (13). In conclusion, these review articles provide a comprehensive overview of T cell biology and offer ideas for further implications in human diseases.

REFERENCES

- Cooper MD, Peterson RD, Good RA. Delineation of the thymic and bursal lymphoid systems in the chicken. *Nature* 1965;205:143-146.
 PUBMED | CROSSREF
- Zinkernagel RM, Doherty PC. Restriction of *in vitro* T cell-mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semiallogeneic system. *Nature* 1974;248:701-702.
 PUBMED | CROSSREF
- 3. Chung Y, Lee SW, Kim WU. Golden age of immunotherapy: challenges and opportunities. *Immune Netw* 2022;22:e1.

PUBMED | CROSSREF

- Lee SW, Lee GW, Kim HO, Cho JH. Shaping heterogeneity of naive CD8⁺ T cell pools. *Immune Netw* 2023;23:e2.
 CROSSREF
- Kim HR, Park JS, Soh WC, Kim NY, Moon HY, Lee JS, Jun CD. T cell microvilli: finger-shaped external structures linked to the fate of T cells. *Immune Netw* 2023;23:e3.
- Lee GR. Molecular mechanisms of T helper cell differentiation and functional specialization. *Immune Netw* 2023;23:e4.
 CROSSREF
- Friedman MJ, Lee H, Lee JY, Oh S. Transcriptional and epigenetic regulation of context-dependent plasticity in T-helper lineages. *Immune Netw* 2023;23:e5.
- Yoon JH, Do JS, Velankanni P, Lee CG, Kwon HK. Gut microbial metabolites on host immune responses in health and disease. *Immune Netw* 2023;23:e6.
 CROSSREF
- Lee C, Lee H, Park JC, Im SH. Microbial components and effector molecules in T helper cell differentiation and function. *Immune Netw* 2023;23:e7.
 CROSSREF

- Koh JY, Kim DU, Moon BH, Shin EC. Human CD8⁺ T-cell populations that express natural killer receptors. Immune Netw 2023;23:e8.
 CROSSREF
- 11. Amitrano AM, Kim M. Metabolic challenges in anticancer CD8 T cell functions. *Immune Netw* 2023;23:e9. CROSSREF
- Choi H, Kim Y, Jung YW. The function of memory CD8+ T cells in immunotherapy for human diseases. *Immune Netw* 2023;23:e10. CROSSREF
- 13. Seok J, Cho SD, Seo SJ, Park SH. Roles of virtual memory T cells in diseases. *Immune Netw* 2023;23:e11. CROSSREF